

Universal Spectral Correlation between Hamiltonians with Disorder

E. BRÉZIN¹ AND A. ZEE²

¹*Laboratoire de Physique Théorique
École Normale Supérieure
24 Rue Lhomond
75231 Paris, France*

²*Institute for Theoretical Physics
University of California
Santa Barbara, CA 93106-4030 USA*

ABSTRACT: We study the correlation between the energy spectra of two disordered Hamiltonians of the form $H_a = H_{0a} + s_a \varphi$ ($a=1,2$) with H_{0a} and φ drawn from random distributions. We calculate this correlation function explicitly and show that it has a simple universal form for a broad class of random distributions.

In a series of recent papers [BZ1, BZ2, BZ3] (hereafter to be referred to as I, II, and III respectively) we studied the theory of random matrices [WIG,POR,MEH] and discovered a remarkably simple universality in the correlation between energy eigenvalues of random Hamiltonians. A brief review and summary of our work may be found in III. In particular, we find that for a large class of Hamiltonians, the smoothed connected correlation (to be defined below) between the density of energy eigenvalues at energies μ and ν has the universal form

$$\rho_c^{\text{smooth}}(\mu, \nu) = \frac{-1}{2(N\pi a)^2} \frac{1}{(x-y)^2} \left\{ \frac{(1-xy)}{[(1-x^2)(1-y^2)]^{1/2}} \right\}. \quad (\text{universal})$$

(where we have centered the spectrum at the origin.) Here a denotes the endpoint of the spectrum and we have introduced the scaled variables $x = \mu/a$ and $y = \nu/a$. We see that up to a factor of $1/a^2$, ρ_c^{smooth} is equal to a universal function of x and y . The only dependence on the random distribution of Hamiltonians appears through a . This

result is all the more remarkable since, by way of contrast, the density $\rho(\mu)$ is completely non-universal. This universality has also been obtained recently by Beenakker [bee] using another method. For μ and ν far from the endpoints of the spectrum, we see that in (universal) the factor in the curly bracket is essentially equal to one, and so the form the universal correlation function basically says that there is an algebraic long-distance correlation falling like $1/(x - y)^2$. In III, we developed an efficient diagrammatic method to calculate the density of eigenvalues and the correlation between them.

Meanwhile, in the recent literature Altschuler and collaborators [SSA] have considered the correlation between the energy spectra of two Hamiltonians H_1 and H_2 . In particular, Simons and Altschuler considered the problem of a single electron moving in a ring threaded by a magnetic flux and with the electron scattering on impurities in the ring. The magnetic flux is then changed to some other value. Here $H_a = H_{0a} + \varphi$ for $a = 1, 2$, where H_{0a} are deterministic, representing the interaction of the electron with the magnetic flux, while φ is random, representing the interaction of the electron with the impurities. The correlation between the spectra of H_1 and H_2 is of interest in the physics of mesoscopic systems. In III, we determined this correlation for a certain class of distribution of φ .

More generally, we may consider the correlation between the spectra of two Hamiltonians $H_a = H_{0a} + s_a\varphi$ for $a = 1, 2$, where s_a represent the strength of the disorder. Simons, Altschuler, and Lee [SSA] and more recently Narayan and Shastry [shastry] have considered this problem for the simpler case of $H_{01} = H_{02} \equiv H_0$. In this paper, we would like to demonstrate the power of the diagrammatic approach discussed in III by calculating the correlation between the spectra of H_1 and H_2 for the more general case in which $H_{01} \neq H_{02}$. Furthermore, instead of obtaining an implicit integral representation for the correlation, we will determine it explicitly.

It may be useful to clarify the difference [diff] between our work and the work of Altschuler et al. They focus [SSA,shastry] on the correlation function $\rho_c(\mu, \nu)$ for $\mu - \nu$ small, of order $1/N$, (that is, for “distances” much smaller than the width of the spectrum), and indeed often for μ and ν both small. In III and in this work (and in the recent work of Beenakker [bee]) we look at the correlation function for $\mu - \nu$ of order N^0 . In other words, we look at long distance correlation, while Altschuler and collaborators focus on short distance correlation. In fact, in I, by determining the large-index behavior of the relevant orthogonal polynomials, we were able to calculate explicitly the *complete* correlation function over all “distance” scales, from short distances through intermediate distances to long distances. The correlation function discussed in the literature appears as

a limiting case of the correlation function found in I. The work in I thus provides a bridge between the short distance discussion of Altschuler et al and the long distance discussion we gave in II and III.

Let us begin by recalling some basic definitions and formulas. Consider an ensemble of N by N hermitean matrices φ defined by a probability distribution $P(\varphi)$ and normalized by $\int d\varphi P(\varphi) = 1$. The large N limit will always be understood. Define as usual the Green's functions

$$G(z) \equiv \left\langle \frac{1}{N} \text{tr} \frac{1}{z - \varphi} \right\rangle \quad (1.3)$$

$$G(z, w)_c \equiv \left\langle \frac{1}{N} \text{tr} \frac{1}{z - \varphi} \frac{1}{N} \text{tr} \frac{1}{w - \varphi} \right\rangle_c \quad (1.4)$$

where $\langle 0(\varphi) \rangle \equiv \int d\varphi 0(\varphi) P(\varphi)$ and the subscript c indicates the connected Green's function. The density of eigenvalues is then given by

$$\rho(\mu) = \left\langle \frac{1}{N} \text{tr} \delta(\mu - \varphi) \right\rangle = \frac{-1}{\pi} \text{Im} G(\mu + i\epsilon) \quad (1.6)$$

and the correlation between eigenvalues, by

$$\begin{aligned} \rho_c(\mu, \nu) &= \left\langle \frac{1}{N} \text{tr} \delta(\mu - \varphi) \frac{1}{N} \text{tr} \delta(\nu - \varphi) \right\rangle_c \\ &= (-1/4\pi^2)(G_c(++) + G_c(--)) - G_c(+-) - G_c(-+) \end{aligned} \quad (1.7)$$

with the obvious notation $G_c(\pm, \pm) \equiv G_c(\mu \pm i\epsilon, \nu \pm i\delta)$ (signs uncorrelated). It is worth remarking that there are two limits involved: $N \rightarrow \infty$ and $\epsilon, \delta \rightarrow 0$. As explained in III, we obtain the full correlation function (from which we can read off the correlation at all distances) by letting $\epsilon, \delta \rightarrow 0$ first (the limit we took in I), and the smoothed correlation function by taking $N \rightarrow \infty$ first. In this paper, we will look at the smoothed correlation.

For applications to disordered system φ is to be thought of as the Hamiltonian. Its eigenvalues then describe the energy levels of the system. In some applications, φ is related to the transmission matrix [PICH]. From the density of and correlation between its eigenvalues we learn about the fluctuation of the conductance in disordered metals.

In this work, we wish to study the correlation between the spectra of two different Hamiltonians H_1 and H_2 . Evidently, we can determine this quantity by studying the generalized (connected) Green's function

$$\mathcal{G}(z, w)_c \equiv \left\langle \frac{1}{N} \text{tr} \frac{1}{z - \varphi_1} \frac{1}{N} \text{tr} \frac{1}{w - \varphi_2} \right\rangle_c \quad (\text{cross}G)$$

where to conform to field theoretic notation we have renamed H_a as φ_a , two “matrix scalar fields.” Note that our starting point appears to correspond to one of the concluding results of Ref. shastri. (Cf. eq. (16) there.)

For ease of presentation we will begin with the Gaussian distribution

$$P(H_{01}, H_{02}, \varphi) = \frac{1}{Z} e^{-\frac{N}{2} \text{tr}[m_1^2 H_{01}^2 + m_2^2 H_{02}^2 + m^2 \varphi^2 + \mu^2 (H_{01} - H_{02})^2]}. \quad (\text{distribution})$$

Note that the factors of N are chosen in our definitions such that in the large N limit the density of eigenvalues has a finite (*i.e.*, of order N^0) width. For $\mu^2 \rightarrow \infty$, we recover the case $H_{01} = H_{02}$ considered earlier [SSA, shastri, BZ3]. In the opposite limit $\mu^2 = 0$, the two Hamiltonians H_{01} and H_{02} are completely independent. In evaluating $\mathcal{G}(z, w)_c$, we see that, since the integrand depends only on φ_a , after using the definition $\varphi_a = H_{0a} + s_a \varphi$ to eliminate H_{0a} and integrating over φ , we can effectively use $P(\varphi) = \frac{1}{Z} e^{-\frac{N}{2} \text{tr}(\varphi_a M_{ab}^2 \varphi_b)}$ with the 2-by-2 symmetric matrix M^2 defined by

$$M_{11}^2 = (m_1^2 + \mu^2) - \mathcal{M}^{-2}[(m_1^2 + \mu^2)s_1 - \mu^2 s_2]^2, \quad (m1)$$

and similarly for M_{22}^2 , and

$$M_{12}^2 = -\mu^2 - \mathcal{M}^{-2}[(m_1^2 + \mu^2)s_1 - \mu^2 s_2][(m_2^2 + \mu^2)s_2 - \mu^2 s_1], \quad (m2)$$

with

$$\mathcal{M}^2 \equiv m^2 + \sum ((m_a^2 + \mu^2)s_a^2 - 2s_1 s_2 \mu^2). \quad (m3)$$

Let us now expand

$$\mathcal{G}_c(z, w) = \frac{1}{N^2} \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{1}{z^{m+1} w^{n+1}} \langle \text{tr} \varphi_1^m(t) \text{tr} \varphi_2^n(0) \rangle_c \quad (\text{expand})$$

Diagrammatically, we may borrow the terminology of large N QCD [thoo] and describe the expression for $\mathcal{G}_c(z, w)$ as two separate quark loops, of type z and type w respectively, interacting by emitting and absorbing gluons. A quark of type z can only emit and absorb gluons of “type 1” and a quark of type w and only emit and absorb gluons of “type 2.” With a Gaussian distribution for φ_a , $a = 1, 2$, we can readily “Wick-contract” (expand). The inverse gluon propagator is given by

$$\langle \varphi_{aij} \varphi_{bkl}(0) \rangle = \frac{1}{N} \delta_{il} \delta_{jk} \sigma_{ab}^2 \quad (\text{glueprop})$$

where the 2-by-2 matrix $\sigma^2 \equiv M^{-2}$. (For the sake of notational simplicity, we will define $\sigma^2 = (\sigma^2)_{11}$, $\tau^2 = (\sigma^2)_{22}$, and $\rho^2 = (\sigma^2)_{12}$.) Thus, the gluon is represented by a double line

while a quark is represented by a single line. This convention greatly facilitates counting the powers of N .

Let us begin by ignoring contractions within the same trace (in which case m and n are required to be equal). In the large N limit, the dominant graphs are given essentially by “ladder graphs” (with one crossing) which immediately sum to

$$N^2 \mathcal{G}_c(z, w) = \frac{\rho^2}{(zw)^2} \frac{1}{(1 - \frac{\rho^2}{zw})^2} \quad (\text{ladder})$$

We next include Wick-contractions within the same trace in $\langle \text{tr} \varphi_1^m \text{tr} \varphi_2^n \rangle$. Graphically these contractions correspond to decorating the ladder graphs by vertex and self energy corrections. Summing the vertex corrections, we see that we have to multiply the expression in (ladder) by two factors, the factor $(1 - \frac{\sigma^2}{z^2})^{-2}$ and a similar factor with z and σ replaced by w and τ respectively. Finally, summing the self energy corrections we see that the bare quark propagator $1/z$ gets dressed to

$$G(z, \sigma) = \frac{1}{2\sigma^2} (z - \sqrt{z^2 - 4\sigma^2}) \quad (\text{green})$$

(and similarly for $1/w$ of course with σ replaced by τ .) Note that the bare propagators are recovered in the limit $\sigma, \tau \rightarrow 0$. Putting these various factors together, we obtain the remarkably compact result

$$N^2 \mathcal{G}_c(z, w) = \frac{\rho^2}{(1 - \rho^2 G(z, \sigma) G(w, \tau))^2} \left[\frac{G^2(z, \sigma)}{1 - \sigma^2 G^2(z, \sigma)} \right] \left[\frac{G^2(w, \tau)}{1 - \tau^2 G^2(w, \tau)} \right] \quad (\text{connected2point})$$

As indicated in III, we find it convenient to introduce angular variables: $\sin \theta \equiv \mu/2\sigma$ and $\sin \phi \equiv \nu/2\tau$. As θ and ϕ vary from $-\pi/2$ to $\pi/2$, μ and ν vary over the width of their respective spectra. These angular variables are the natural scaling variables to use in this class of problems dealing with eigenvalues of random matrices. From (green), we have

$$G(\mu + i\epsilon, \sigma) = -i\eta e^{i\eta\theta} / \sigma \quad (\text{cutG1})$$

where $\eta = \text{the sign of } \epsilon$ and

$$G(\nu + i\delta, \tau) = -i\xi e^{i\xi\phi} / \tau \quad (\text{cutG2})$$

with $\xi = \text{the sign of } \delta$. A straightforward computation of the appropriate generalization of (1.7) then yields

$$\begin{aligned} & -16\pi^2 N^2 \rho_c(\mu, \nu) \\ &= \frac{1}{\sigma\tau \cos \theta \cos \phi} \left\{ \frac{1 + \text{ch } u \cos(\theta + \phi)}{[\text{chu} + \cos(\theta + \phi)]^2} + \frac{1 - \text{ch } u \cos(\theta - \phi)}{[\text{chu} - \cos(\theta - \phi)]^2} \right\}. \end{aligned} \quad (\text{correlation})$$

Here we have defined

$$u \equiv \log(\sigma\tau/\rho^2) = \frac{1}{2} \log[M_{11}^2 M_{22}^2 / (M_{12}^2)^2], \quad (\text{define } u)$$

a rather involved function of the parameters m_1^2 , m_2^2 , m^2 , and μ^2 that appear in the distribution (distribution). Perhaps remarkably, aside from an overall factor of $\sigma\tau$ the correlation function, when expressed in terms of the appropriately scaled variables θ and ϕ depends on the probability distribution only through this particular combination. This represents already a form of universality or scaling.

The reader familiar with III would recognize that this is the same expression we obtained there (see eq. (2.18) in Ref. BZ3) for an apparently totally different problem, with a totally different definition of u . In III we studied the time-dependent correlation between the eigenvalues of time-dependent matrices $\varphi(t)$ taken from a probability distribution defined by

$$P(\varphi) = \frac{1}{Z''} \exp - \int_{-T}^T dt \operatorname{Tr} \left[\varphi K\left(\frac{d}{dt}\right) \varphi + V(\varphi) \right] \quad (\text{time } dep)$$

with K any reasonable function and with $T \rightarrow \infty$. Here “time” may correspond to some external parameter we are allowed to vary. For this problem, we defined $e^{-u(t)}$ to be $\int \frac{d\omega}{2\pi} e^{i\omega t} (1/K(\omega))$. For the problem considered in this paper, in contrast, e^{-u} is defined simply in terms of the widths of the distributions from which various random Hamiltonians are drawn. Evidently, we have in this way found a “mapping” between two apparently unrelated problems involving disordered Hamiltonians. A particularly simple case occurs when $K(\frac{d}{dt}) = -\frac{d^2}{dt^2}$ in which case $u(t)$ is essentially time. It is in this sense that loosely, one may speak of u as defined in (define u) as “time,” even though there is no notion of time as such in the problem studied in this paper.

Various limits can now be studied. In the large “time” limit $u \rightarrow \infty$, we obtain

$$4\pi^2 N^2 \rho_c(\mu, \nu) \rightarrow e^{-u} \tan \theta \tan \phi \quad (\text{longtime})$$

For $\theta = \phi$ (note that this does not mean $\mu = \nu$ when σ is not equal to τ) and small “time” u we find

$$8\pi^2 N^2 \rho_c(\mu, \nu) = \frac{1}{\sigma\tau \cos^2 \theta} \frac{1}{u^2} + \dots \quad (\text{samepoint})$$

For $\theta \neq \phi$ and $u \ll \theta - \phi$ we obtain

$$-16\pi^2 N^2 \rho_c(\mu, \nu) = \frac{1}{\sigma\tau \cos \theta \cos \phi} \left\{ \frac{1}{1 + \cos(\theta + \phi)} \left[1 - \frac{1 - \frac{1}{2} \cos(\theta + \phi)}{1 + \cos(\theta + \phi)} u^2 \right] + (\phi \rightarrow -\phi + \pi) \right\} \quad (\text{smalltime})$$

We can also study various regions of parameter space, of course. Consider (A) $\mu \rightarrow \infty$ so that $H_{01} = H_{02} = H_0$ in which case it is convenient to define $m_0^2 = m_1^2 + m_2^2$. Case (A1): with $m_0 \rightarrow \infty$ (so that $H_0 = 0$) and $s_1 = s_2$ we recover the universality (universal) found in I and in Ref. bee. Case (A2): with $m_0 \gg m$ so that the disorder is large, we find that $e^u = \sigma\tau/\rho^2 = 1$ and thus the same universal correlation as in case (A1) suitably scaled by an overall factor of $\sigma\tau$, but here there is no requirement on s_a . Case (A3): with s_2 equal to zero, so that we are studying the correlation between the spectra of a “bare” Hamiltonian H_0 and a Hamiltonian disturbed by an external perturbation $H_0 + s_1\varphi$, we find that $e^u = 1 + m_0^2 s_1^2 / m^2$. Note that no general statement on how “time” t maps onto the strength s_1 of the external perturbation can be made, but in the simplest case of identifying “time” mentioned above we have $t \propto \log s_1$ for large strength of the perturbation.

Another interesting class (B) is defined by the limit $\mu \rightarrow 0$ so that H_{01} and H_{02} are not correlated at all. After some drastic simplification, we find $\sigma^2 = m_1^{-2} + s_1^2 m^{-2}$, $\tau^2 = m_2^{-2} + s_2^2 m^{-2}$, and $\rho^2 = s_1 s_2 m^{-2}$. As a check on the formalism, we can consider some special cases. For instance, with $m_a \rightarrow \infty, a = 1, 2$, H_{01} and H_{02} are set to zero, in which case we recover essentially our previous result (universal). As another example, consider the analog of case (A3): with $s_2 = 0$ we obtain $u \rightarrow \infty$ and as expected $\rho_c = 0$ since now H_1 and H_2 are no longer “linked” by φ .

All these results are derived with the Gaussian distribution (distribution). We now remark on how these results may hold for more general distribution. In III we identified two classes of random matrix distribution which we refer to as the Wigner class and the trace class. In the Wigner class, we consider an ensemble of matrices whose matrix elements are independently distributed according to some probability distribution (the same distribution for all the elements.) Let us focus on the example in which the probability of the distribution matrix element φ_{ij} is given by

$$P(\varphi_{ij}) \propto e^{-N^2(|\varphi_{ij}|^2 - \frac{v^2}{N^2})^2}. \quad (wigner)$$

(Here φ represents any of the matrices in (distribution) generically.) It is easy to see that in the Feynman diagrams involving the quartic interaction $\sim N^2 |\varphi_{ij}|^4$ the indices i and j are tied together so to speak and thus these graphs are suppressed by a power of N^{-1} relative to the graphs in which this interaction vertex does not appear. Reasoning along this line, we see immediately that the results of this paper have a generality far beyond the Gaussian distribution. Our results are universal. This universality is much harder to

prove for the so-called trace class in which the generic random matrix φ is taken from a distribution of the form $P(\varphi) = \frac{1}{Z} e^{-N \text{tr} V(\varphi)}$. We conjecture however that our results may also hold for random Hamiltonians defined with this class of probability distribution. In II, we had shown that the universal correlation in (universal) indeed holds for this trace class and for a much broader class generalizing the trace class. In III, we also calculated the correlation when H_{01} and H_{02} are set to equal deterministic, rather than random, Hamiltonians W_1 and W_2 say. This case may be considered in the present context by adding to the logarithm of the distribution in (distribution) terms like $-\alpha_a^2 (H_{0a} - W_a)^2$ and taking $\alpha_a \rightarrow \infty$. The gluons φ_1 and φ_2 are now coupled also to external sources.

In conclusion, we have extended the universality in (universal) to a much broader class of problems involving random Hamiltonians. Expressed in terms of suitably scaled variables, the correlation between energy eigenvalues is universal, certainly in the Wigner class, and most likely in the trace class as well. The case $H_{01} = H_{02}$ discussed previously in the literature can be recovered as a special case within our discussion.

ACKNOWLEDGEMENT

This work was supported in part by the National Science Foundation under Grant No. PHY89-04035 and the Institut Universitaire de France.

REFERENCES

- BZ1. E. Brézin and A. Zee, *Nucl. Phys.* **402(FS)**, 613 (1993).
- BZ2. E. Brézin and A. Zee, *Compt. Rend. Acad. Sci.*, 317, 735, 1993.
- diff. One of us (AZ) thanks B. Altschuler for a discussion clarifying this difference.
- BZ3. E. Brézin and A. Zee, “Correlation functions in disordered systems,” SBITP-preprint 1993.

- WIG. E. Wigner, *Can. Math. Congr. Proc.* p.174 (University of Toronto Press) and other papers reprinted in Porter, op. cit.
- POR. C.E. Porter, *Statistical Theories of Spectra: Fluctuations* (Academic Press, New York, 1965).
- MEH. M.L. Mehta, *Random Matrices* (Academic Press, New York, 1991).
- shastry. O. Narayan and B. S. Shastri, *Phys. Rev. Lett.* **71**, 2106 (1993).
- bee. C.W.J. Beenakker, Institut Lorentz preprint (1993) cond-mat/9310010; for earlier work, see C.W.J. Beenakker, *Phys. Rev. Lett.* **70**, 1155 (1993); *Phys. Rev.* **B47**, 15763 (1993).
- SSA. B.D. Simons and B.L. Altschuler, *Phys. Rev. Lett.* **70**, 4063 (1993); B. D. Simons, P. A. Lee, and B. L. Altschuler, *Phys. Rev. Lett.* **70**, 4122 (1993).
- PICH. J.-L. Pichard, in *Quantum Coherence in Mesoscopic Systems* (NATO ASI series, Plenum, New York 1991) ed. by B. Kramer.
- thoo. G. 't Hooft, *Nucl. Phys.* **B 72**, 461 (1974).